

PiXL KnowIT!

GCSE Chemistry

AQA Topic – Energy Changes

© Copyright The PiXL Club Ltd, 2017

This resource is strictly for the use of member schools for as long as they remain members of The PiXL Club. It may not be copied, sold nor transferred to a third party or used by the school after membership ceases. Until such time it may be freely used within the member school.

All opinions and contributions are those of the authors. The contents of this resource are not connected with nor endorsed by any other company, organisation or institution.

Exothermic and endothermic reactions

- Energy changes during exothermic and endothermic reactions
- Reaction profiles
- The energy change of reactions (HT only)

Chemical cells and fuel cells (Chemistry only)

- Cells and batteries
- Fuel cells



LearnIT! KnowIT!

Exothermic and Endothermic reactions

- Exothermic reactions
- Endothermic reactions



Exothermic and endothermic reactions part 1 – Exothermic reactions

Energy is **conserved** in chemical reactions. The amount of energy in the Universe **at the end** of a chemical reaction **is the same as before** the reaction takes place.



In the above reaction energy is **released**, it gets hotter.

An **exothermic reaction** is one that **transfers energy** to the surroundings so the temperature of the surroundings **increases** – “it gets hotter”.

The two HCl molecules made will not hold as much energy as the H₂ and Cl₂ molecules at the start, so the spare energy is released as heat.

Exothermic and endothermic reactions part 1 – Exothermic reactions

There are a number of common exothermic reactions, they include:

Combustion




Know all three of
these examples of
exothermic
reactions

Oxidation



Everyday uses of
exothermic reactions
include –
Self-heating cans
Hand warmers

Neutralisation



Know both of
these uses for
exothermic
reactions

Exothermic and endothermic reactions part 1 – Endothermic reactions

We have already learnt that energy is **conserved** in chemical reactions.



In the above reaction, energy is **taken in**- it gets **colder**.

An **endothermic** reaction is one that takes energy from the surroundings so the temperature of the surroundings **decreases** – “it gets colder”.

The sodium ethanoate, carbon dioxide and water molecules made will hold **more** energy than the ethanoic acid and sodium carbonate molecules at the start, so the energy needed is **taken in** as heat.

Other examples of endothermic reactions are

- **Thermal decomposition**
- **Sports injury packs**

Know all three of these examples of endothermic reactions

QuestionIT!

Energy Changes Part 1

- Exothermic reactions
- Endothermic reactions



1. How would you know if an exothermic reaction had occurred?
2. How would you know if an endothermic reaction had occurred?
3. Below is a table of results for four reactions, the temperatures before and after the reactions are also given.

Reaction	Temperature at start $^{\circ}\text{C}$	Temperature at end $^{\circ}\text{C}$
A	22	28
B	20	20
C	21	12
D	25	25

- a/ Which reaction is endothermic? Explain how you know this.
- b/ Which reaction is exothermic? Explain how you know this.

AnswerIT!

Energy Changes Part 1

- Exothermic reactions
- Endothermic reactions



1. How would you know if an exothermic reaction had occurred?

The reaction would give out heat/get warmer/ temperature increase.

2. How would you know if an endothermic reaction had occurred?

The reaction would take in heat/get colder/ temperature decrease.

3. Below is a table of results for four reactions, the temperatures before and after the reactions are also given.

Reaction	Temperature at start $^{\circ}\text{C}$	Temperature at end $^{\circ}\text{C}$
A	22	28
B	20	20
C	21	12
D	25	25

a/ Which reaction is endothermic? Explain how you know this.

Reaction C, it gets colder/temperature falls.

b/ Which reaction is exothermic? Explain how you know this.


Reaction A, it gets warmer/temperature increases.

Energy Changes

Part 2

-

Chemical reactions can only occur when reacting particles collide with each other **with** sufficient energy.



The **minimum** amount of energy that particles must have to react is called the **activation energy**

You have given a reaction its **activation energy** when you have used a lit spill to light a Bunsen burner. Without the **activation energy** from the lit spill the methane gas and oxygen in the air will not **combust** and release the heat energy.

When we look at this reaction we see the following.




You will be expected to balance this equation.

Energy Changes part 2 – Reaction profiles





We know this reaction is **exothermic**, this means energy is released. So the CH_4 and 2O_2 , the reactants, must have more energy than the products, CO_2 and $2\text{H}_2\text{O}$


We can show this as a **reaction profile**. On it we need to include the formulae or names of the products and reactants. We also need to show the relative energies of the reactants and products



More information needs to be included in the reaction profile. This will show the **activation energy** of the reaction. It is shown by a curved line rising above the reactants energy.



We can now see the overall change in energy within the reaction.




The **products** have **less energy** than the **reactants**. This will have been lost as heat as the reaction is exothermic.

We saw earlier that the following reaction was **endothermic**:



What would the **reaction profile** look like for this reaction?



We can see that the **products** have **more energy** than the **reactants**. This will have been taken in as heat energy it feels colder.

QuestionIT!

Energy Changes part 2


- Activation energy
- Reaction profiles



1. What is meant by the term activation energy?

2. On the reaction profile below what is shown by the letters?

A
B
C
D



3. What two things are needed for a chemical reaction to occur?

4. What is an exothermic reaction?

5. What is an endothermic reaction?

AnswerIT!

Energy Changes part 2

- Activation energy
- Reaction profiles




1. What is meant by the term activation energy?

The minimum amount of energy that particles must have to react.

2. On the reaction profile below what is shown by the letters?

- A Reactants
- B Products
- C Activation energy
- D Overall energy change/that the reaction is exothermic



3. What two things are needed for a chemical reaction to occur?

Reacting particles collide with each other and with sufficient energy.

4. What is an exothermic reaction?

Heat energy given out/energy lost to the surroundings


5. What is an endothermic reaction?

Heat energy taken in/energy taken in from the surroundings

For the reaction of methane with oxygen we can write out the balanced symbol equation:



We can draw out the bonds between the atoms e.g.



Each line represents a bond, two lines represent a **double bond** e.g. in the oxygen molecule.

During a **chemical reaction**:

- Energy must be supplied to **break bonds** in the reactants
- Energy is released to **form bonds** in the products.


The energy needed to break bonds and the energy released when bonds are formed can be calculated from **bond energies**.

Bond	Bond Energy kJ/mol
C-H	411
O=O	494
C=O	799
O-H	459

This means that 411kJ/mol of energy needs to be put in to break the carbon-hydrogen bond.
It also means that 459 kJ/mol is given out when the oxygen hydrogen bond is made in water.

The **difference** between the sum of the energy needed to break bonds in the reactants and the sum of the energy released when bonds in the products are formed is the **overall energy change** of the reaction.

Worked example



For the reactants	For the products
There are four C-H bonds so 4 x 411kJ/mol = 1,644kJ/mol	There are two C=O bonds so 2 x 799kJ/mol = 1,598kJ/mol
There are two O=O bonds so 2 x 494kJ/mol = 988kJ/mol	There are four O-H bonds so 4 x 459kJ/mol = 1,836kJ/mol
The sum of these is the energy supplied to break the bonds in the reactants it is 1,644 + 988 = 2,632kJ/mol	The sum of these is the energy released when bonds in the products are formed it is 1,598 + 1,836 = 3,434kJ/mol

We already know that the **difference** between the sum of the energy needed to break bonds in the reactants and the sum of the energy released when bonds in the products are formed is the **overall energy change** of the reaction.

This means that

$$\begin{aligned} \text{Overall energy change} &= \text{energy needed to break the bonds} - \text{energy released as bonds are made} \\ &= 2,632\text{kJ/mol} - 3,434\text{kJ/mol} \end{aligned}$$

$$\text{Overall energy change} = -802\text{kJ/mol}$$

This is an **exothermic** reaction, so the sum of the difference between the calculations is **negative**. For an **endothermic** reaction it would be **positive**.

Students should be able to calculate the energy transferred in chemical reactions using bond energies supplied.

Know these two definitions- they are often asked for in the exam.

In an **exothermic** reaction, the energy released from forming **new bonds is greater** than the energy needed to break existing bonds



In an **endothermic** reaction, the energy needed to **break existing bonds is greater** than the energy released from forming new bonds



QuestionIT!

The energy change
of reactions
(Higher Tier
only)



- 1. Which process is exothermic, bond breaking or bond making?**
- 2. Explain your answer to question 1.**
- 3. How do we calculate the overall energy change of a reaction?**

4. The bond energy between a hydrogen and a nitrogen atom is 386 kJ/mol, the bond energy between two hydrogen atoms is 432 kJ/mol and the bond energy between two nitrogen atoms is 942 kJ/mol. Using these bond energies, calculate the overall energy change for the following reaction.



5. Is the reaction exothermic or endothermic? Explain your answer.

AnswerIT!

The energy change
of reactions
(Higher Tier
only)



1. Which process is exothermic, bond breaking or bond making?

Bond making

2. Explain your answer to question 1

Energy is released/temperature increases

3. How do we calculate the overall energy change of a reaction?

It is the difference between the sum of the energy needed to break bonds in the reactants and the sum of the energy released when bonds in the products are formed.

4. The bond energy between a hydrogen and a nitrogen atom is 386 kJ/mol, the bond energy between two hydrogen atoms is 432 kJ/mol and the bond energy between two nitrogen atoms is 942 kJ/mol. Using these bond energies calculate the overall energy change for the following reaction.



For the reactants

There are three nitrogen to hydrogen bonds in each molecule and there are two molecules so

$$386\text{kJ/mol} \times 6 = \mathbf{2,316\text{kJ/mol}}$$

For the products

There are three hydrogen to hydrogen bonds (432 kJ/mol) and one nitrogen to nitrogen triple bond so

$$3 \times 432\text{kJ/mol} + 942\text{kJ/mol} = \mathbf{2,238\text{kJ/mol}}$$

Overall energy change	=	energy needed to break the bonds	–	energy released as bonds are made
		2,316 kJ/mol	-	2,238 kJ/mol

$$\text{Overall energy change} = +78\text{kJ/mol}$$

5. Is the reaction exothermic or endothermic? Explain your answer.

The reaction is endothermic as the overall energy change is positive.

LearnIT! KnowIT!

Chemical Cells And Fuel Cells (Chemistry Only)



- Cells And Batteries
- Fuel Cells

Cells contain chemicals which react to produce electricity.



The **voltage** produced by a cell is dependent upon a number of factors including the type of **electrode** and **electrolyte**.

A simple **cell** can be made by connecting two different metals in contact with an **electrolyte**.

Batteries consist of two or more **cells** connected together in **series** to provide a greater voltage.

In **non-rechargeable cells** and batteries the chemical reactions stop when one of the reactants has been used up. **Alkaline batteries** are **non-rechargeable**.

Rechargeable cells and **batteries** can be recharged because the chemical reactions are reversed when an external electrical current is supplied.



Fuel cells are supplied by an external source of fuel (e.g. hydrogen) and oxygen or air. The fuel is **oxidised electrochemically** within the fuel cell to produce a **potential difference** (a voltage).

The overall reaction in a hydrogen fuel cell involves the oxidation of hydrogen to produce water.

hydrogen + oxygen → water



(Higher Tier only)

There are two electrodes in the hydrogen fuel cell.


At the **cathode** (+ charged)



At the **anode** (- charged)



Hydrogen fuel cells offer a potential alternative to rechargeable cells and batteries.



The half equations are only needed for the higher tier, however they are very important and worth revising carefully

QuestionIT!

Chemical Cells And Fuel Cells (Chemistry Only)

- Cells And Batteries
- Fuel Cells



1. Give two factors which may affect the voltage given out by a battery.

2. Here is a reactivity series of metals. The most reactive is first, the least reactive is last: Magnesium Zinc Tin Copper

Which two metals would you use to make a battery which had the highest voltage? Explain why.

3. Why do non-renewable batteries stop producing a voltage after a certain time?

4. How are rechargeable batteries recharged?

- 1. What chemical is the fuel in a fuel cell ?**
- 2. What happens to this fuel inside the fuel cell to produce a potential difference?**
- 3. Write the overall balanced symbol equation for the reaction in a fuel cell.**
- 4. Write the half equation for the reaction that happens at the cathode in a fuel cell.**
- 5. Write the half equation for the reaction that happens at the anode in a fuel cell.**

AnswerIT!

Chemical Cells And Fuel Cells (Chemistry Only)



- Cells And Batteries
- Fuel Cells

1. Give two factors which may affect the voltage given out by a battery.

The type of electrode and the electrolyte

2. Here is a reactivity series of metals. The most reactive is first, the least reactive is last. Magnesium Zinc Tin Copper

Which two metals would you use to make a battery which gave off the most voltage? Explain why

Magnesium and copper as largest difference in reactivity.

3. Why do non-renewable batteries stop producing a voltage after a certain time?

The chemical reactions stop when one of the reactants has been used up.

4. How are rechargeable batteries recharged?

The chemical reactions are reversed by an external electrical current.

1. What chemical is the fuel in a fuel cell ?

Hydrogen.

2. What happens to this fuel inside the fuel cell to produce a potential difference?

It is oxidised electrochemically.

3. Write the overall balanced symbol equation for the reaction in a fuel cell.



4. Write the half equation for the reaction that happens at the cathode in a fuel cell.



5. Write the half equation for the reaction that happens at the anode in a fuel cell.

